

AD-A060 884

CALIFORNIA UNIV SAN DIEGO LA JOLLA DEPT OF PHYSICS

F/G 20/10

LOCALITY PRINCIPLE IN WAVE-MECHANICS (WAVE MECHANICS/LOCAL DENS--ETC(U)

OCT 78 W KOHN, A YANIV

N00014-76-C-0050

UNCLASSIFIED


TR-47

NL

1 OF 1
AD
A060 884



END
DATE
FILMED
1-79
DDC



LEVEL II

12 SC

UNCLASSIFIED
Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) DEPARTMENT OF PHYSICS, B-019 UNIVERSITY OF CALIFORNIA SAN DIEGO, LA JOLLA, CA, 92093		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
2b. GROUP			
3. REPORT DATE October 1978			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) TECHNICAL REPORT, (1/1/78-10/25/78) 1 Jan - 25 Oct 78			
5. AUTHOR (Last name, first name) Walter Kohn and Avishay Yaniv			
6. REPORT DATE October 1978		7a. TOTAL NO. OF PAGES 13	7b. NO. OF REFS 7
8. CONTRACT OR GRANT NO. NONR N00014-76-C-00504		9a. ORIGINATOR'S REPORT NUMBER(S) NR17-653	
9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)			
10. DISTRIBUTION STATEMENT APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.			
11. SUPPLEMENTARY NOTES rho		12. SPONSORING MILITARY ACTIVITY OFFICE OF NAVAL RESEARCH DEPARTMENT OF THE NAVY, ARLINGTON, VA. 22217	
13. ABSTRACT This paper proves a locality principle for a wave-mechanical particle governed by the Schroedinger equation. It is shown that $\rho(r, \beta)$, the Laplace transform of the local density of states $n(r, E)$, depends significantly only on the potential $V(r')$ at points r' near r . The effect of changes of $V(r')$ at distant points r' ($ r' - r > d$) on $\rho(r, \beta)$, decay in a Gaussian fashion with a . This result sheds some light on the locality of physical properties of extended systems and provides general support for various local methods of calculation. abs. val. $(r' - r)$			

AD A060884

DDC FILE COPY

073390

DDC
RECEIVED
NOV 6 1978
D

Security Classification

DD FORM 1473 (BACK)
(PAGE 2)

UNCLASSIFIED
Security Classification

ADDITIONAL BY	
BY	Write Section <input checked="" type="checkbox"/>
DO	Self Section <input type="checkbox"/>
UNANNOUNCED <input type="checkbox"/>	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. 200/04 SPECIAL
A	

TECHNICAL REPORT #47 ✓

CONTRACT NONR-N00014-76-C0050

PROJECT NR 017-653

Supervisor: Professor Walter Kohn

DEPARTMENT OF PHYSICS
UNIVERSITY OF CALIFORNIA SAN DIEGO
LA JOLLA, CALIFORNIA

DDC
RECEIVED
NOV 6 1978
D

Reproduction in whole or in part is permitted for any purpose of the United States Government.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

October 1978

78 11 01 068

LOCALITY PRINCIPLE IN WAVE-MECHANICS

(Wave Mechanics/Local Density of States/Local Partition Function/
Cluster Methods/Moment Methods/Path Integral Methods)

W. Kohn and A. Yaniv

Department of Physics, University of California San Diego,

La Jolla, California 92093

ABSTRACT

This paper proves a locality principle for a wave-mechanical particle governed by the Schroedinger equation. It is shown that $\rho(\underline{r}, \beta)$, the Laplace transform of the local density of states $n(\underline{r}, E)$, depends significantly only on the potential $V(\underline{r}')$ at points \underline{r}' near \underline{r} . The effect of changes of $V(\underline{r}')$ at distant points \underline{r}' ($|\underline{r}' - \underline{r}| > d$) on $\rho(\underline{r}, \beta)$, decay in a Gaussian fashion with a . This result sheds some light on the locality of physical properties of extended systems and provides general support for various local methods of calculation.

1. Introduction

This note deals with a particle described by the single particle Hamiltonian

$$H \equiv \frac{1}{2} p^2 + V(\underline{r}) \quad , \quad (1.1)$$

where $V(\underline{r})$ is defined for all \underline{r} inside a large region Ω which eventually becomes infinite. The eigenfunctions and eigenvalues of H , $\psi^V(\underline{r})$ and E^V , are defined by the Schroedinger equation

$$H \psi^V(\underline{r}) = E^V \psi^V(\underline{r}) \quad , \quad (1.2)$$

the boundary condition

$$\psi^V(\underline{r}) = 0 \text{ on boundary of } \Omega, \quad (1.3)$$

and the normalization condition

$$\int |\psi^V(\underline{r})|^2 d\underline{r} = 1 . \quad (1.4)$$

Of course a change of $V(\underline{r})$ in the vicinity of a point \underline{r}_0 in general changes the eigenfunctions $\psi^V(\underline{r})$ at all points \underline{r} . This change can be considerable even if the distance between \underline{r} and \underline{r}_0 becomes large. For example, if we start with $V(\underline{r}) \equiv 0$ and Ω a large box $(-L < x, y, z < +L)$, and subsequently introduce a one dimensional localized barrier

$$V(\underline{r}) \equiv \begin{cases} V_0 & -d < x < d \\ 0 & \text{otherwise} \end{cases} \quad , \quad (1.5)$$

the eigenfunctions $\psi^V(\underline{r})$ are changed by comparable amounts for all values of \underline{r} . This can be regarded as a kind of "action-at-a-distance" of the potential $V(\underline{r})$ on the wavefunctions $\psi^V(\underline{r})$.

Let us now define the quantity

$$\rho(\underline{r}, \beta) \equiv \sum_V |\psi^V(\underline{r})|^2 e^{-\beta E^V}, \quad (1.6)$$

which we call the local partition function (LPF). Its integral over \underline{r} is the conventional partition function,

$$Z(\beta) \equiv \int \rho(\underline{r}, \beta) d\underline{r} = \sum_V e^{-\beta E^V}. \quad (1.7)$$

Also, clearly, the LPF is the Laplace transform of the local density of states

$$n(\underline{r}, E) \equiv \sum_V |\psi^V(\underline{r})|^2 \delta(E - E^V), \quad (1.8)$$

whose spatial integral is the total density of states

$$n(E) \equiv \int n(\underline{r}, E) d\underline{r} = \sum_V \delta(E - E^V). \quad (1.9)$$

Evidently

$$\rho(\underline{r}, \beta) = \int n(\underline{r}, E) e^{-\beta E} dE \quad (1.10)$$

and

$$Z(\beta) = \int n(E) e^{-\beta E} dE. \quad (1.11)$$

We shall show that, in contrast to an individual eigenfunction, $\psi^V(\underline{r})$, the LPF $\rho(\underline{r}, \beta)$ is significantly affected by changes of $V(\underline{r})$ only if these occur at points \underline{r}_0 near \underline{r} . This principle, which will be made more precise below, we call locality principle of wave mechanics.

2. Proof of the Locality Principle

The proof proceeds from the path integral expression for $\rho(\underline{r}, \beta)^{(1)}$. For convenience, we take $\underline{r} = 0$. Then

$$\rho(\beta) \equiv \rho(0, \beta) = \int \exp \left\{ - \int_0^\beta \left(\frac{1}{2} \dot{\underline{r}}^2(u) + V(\underline{r}) \right) du \right\} \mathcal{D}\underline{r}(u), \quad (2.1)$$

where u is an imaginary "time" and $\mathcal{D}\underline{r}(u)$ denotes path integration over

all paths starting and ending at $\underline{r} = 0$.

Let us divide Ω into an interior region

$$\Omega_{\text{int}}: \quad |x|, |y|, |z| \leq a, \quad (2.2)$$

and an exterior region

$$\Omega_{\text{ext}}: \quad |x|, |y|, |z| > a. \quad (2.3)$$

We may then write $\rho(\beta)$ as the sum of two terms,

$$\rho(\beta) = \rho_{\text{int}}(\beta) + \rho_{\text{ext}}(\beta), \quad (2.4)$$

where $\rho_{\text{int}}(\beta)$ comprises all contributions to (2.1) coming from paths entirely inside Ω_{int} , and $\rho_{\text{ext}}(\beta)$ represents the contributions from paths extending outside Ω_{int} . We may note that $\rho_{\text{int}}(\beta)$ is the LPF for a system with potential $V(\underline{r})$ inside Ω_{int} and infinite walls at the boundary of Ω_{int} .

Now consider a comparison potential, $V'(\underline{r})$, identical to $V(\underline{r})$ inside Ω_{int} but different in Ω_{ext} . The LPF for this system can be written as

$$\rho'(\beta) = \rho_{\text{int}}(\beta) + \rho'_{\text{ext}}(\beta), \quad (2.5)$$

where the interior contribution is, by the construction (2.1) the same as for $\rho(\beta)$, Eq. (2.4).

Thus the difference of the two LPFs is given by

$$\Delta\rho(\beta) \equiv \rho'(\beta) - \rho(\beta) = \rho'_{\text{ext}}(\beta) - \rho_{\text{ext}}(\beta). \quad (2.6)$$

Now let us assume that both $V(\underline{r})$ and $V'(\underline{r})$ are bounded below

$$V(\underline{r}), V'(\underline{r}) \geq V_{\text{min}}. \quad (2.7)$$

Then clearly, from (2.1),

$$\begin{aligned} \rho_{\text{ext}}(\beta), \quad \rho'_{\text{ext}}(\beta) &\leq \int \exp \left\{ -\int_0^\beta \left(\frac{1}{2} \dot{\underline{r}}^2(u) + V_{\min} \right) du \right\} \mathcal{D}_{\text{ext}} \underline{r}(u) \\ &= e^{-\beta V_{\min}} \rho_{0,\text{ext}}(\beta), \end{aligned} \quad (2.8)$$

where $\mathcal{D}_{\text{ext}} \underline{r}(u)$ denotes integration over all paths which extend into Ω_{ext} and

$$\rho_{0,\text{ext}}(\beta) \equiv \int \exp \left\{ -\int_0^\beta \frac{1}{2} \dot{\underline{r}}^2(u) du \right\} \mathcal{D}_{\text{ext}} \underline{r}(u) \quad (2.9)$$

corresponds to a free particle, $V(\underline{r}) \equiv 0$. Next let us assume that $V(\underline{r})$ (but not necessarily $V'(\underline{r})$) is bounded above,

$$V(\underline{r}) \leq V_{\max}. \quad (2.10)$$

Then clearly

$$\rho(\beta) \geq e^{-\beta V_{\max}} \rho_0(\beta), \quad (2.11)$$

where $\rho_0(\beta)$ is the local partition function for a free particle,

$$\begin{aligned} \rho_0(\beta) &\equiv \int \exp \left\{ -\int_0^\beta \frac{1}{2} \dot{\underline{r}}^2(u) du \right\} \mathcal{D} \underline{r}(u) \\ &= \left(\frac{1}{2\pi\beta} \right)^{3/2}. \end{aligned} \quad (2.12)$$

Combining Eqs. (2.6), (2.8), (2.11) and (2.12) gives

$$\frac{|\Delta \rho(\beta)|}{\rho(\beta)} \leq 2 e^{\beta(V_{\max} - V_{\min})} \frac{\rho_{0,\text{ext}}(\beta)}{\rho_0(\beta)}. \quad (2.13)$$

Lastly we calculate $\rho_{0,\text{ext}}(\beta)$, the contribution to $\rho_0(\beta)$ of paths extending beyond Ω_{int} . We write

$$\rho_{0,\text{ext}}(\beta) = \rho_0(\beta) - \rho_{0,\text{int}}(\beta) \quad (2.14)$$

and recall that $\rho_{o,int}(\beta)$ is the LPF for free electrons in the box (2.2) with infinite walls. The cosine eigenfunctions and eigenvalues are

$$\psi^v(r) = \frac{1}{a^{3/2}} \cos\left[\frac{\pi(v_1 + \frac{1}{2})}{a} x\right] \cos\left[\frac{\pi(v_2 + \frac{1}{2})}{a} y\right] \cos\left[\frac{\pi(v_3 + \frac{1}{2})}{a} z\right], \quad (2.15)$$

$$E^v = \frac{1}{2} \frac{\pi^2}{a^2} \left[(v_1 + \frac{1}{2})^2 + (v_2 + \frac{1}{2})^2 + (v_3 + \frac{1}{2})^2 \right], \quad (2.16)$$

where $v \equiv (v_1, v_2, v_3)$. Eigenfunctions which are odd in x , y or z do not contribute to the LPF at $r = 0$. Therefore we have

$$\begin{aligned} \rho_{o,int}(\beta) &\equiv a^{-3} \sum_{v_1, v_2, v_3=0}^{\infty} e^{-\frac{\pi^2}{2a^2} \left[(v_1 + \frac{1}{2})^2 + (v_2 + \frac{1}{2})^2 + (v_3 + \frac{1}{2})^2 \right]} \\ &= a^{-3} \left[\sum_{\mu} e^{-\beta \frac{\pi^2}{2a^2} (\mu + \frac{1}{2})^2} \right]^3. \end{aligned} \quad (2.17)$$

Using Poisson's summation formula⁽²⁾ we can rewrite the sum over μ as

$$a^{-1} \sum_{\mu} e^{-\beta \frac{\pi^2}{2a^2} (\mu + \frac{1}{2})^2} = (2\pi\beta)^{-1/2} \left(1 - 2 \sum_{n=1}^{\infty} (-1)^{n+1} e^{-\frac{2n^2 a^2}{\beta}} \right) \quad (2.18)$$

so that

$$\rho_{o,ext}(\beta) = (2\pi\beta)^{-3/2} \left\{ 1 - (1 - 2 \sum_{n=1}^{\infty} (-1)^{n+1} e^{-\frac{2n^2 a^2}{\beta}})^3 \right\}. \quad (2.19)$$

For large a only the leading term in the sum needs to be retained giving

$$\rho_{o,ext}(\beta) \rightarrow (2\pi\beta)^{-3/2} 6 e^{-\frac{2a^2}{\beta}}. \quad (2.20)$$

It can be verified that the right hand side of (2.20) constitutes an upper bound on $\rho_{o,ext}(\beta)$ for all a ,

$$\rho_{0,\text{ext}}(\beta) \leq (2\pi\beta)^{-\frac{3}{2}} 6 e^{-\frac{2a^2}{\beta}}. \quad (2.21)$$

Substitution of (2.12) and (2.21) into (2.13) gives our final result

$$\frac{|\Delta\rho(\beta)|}{\rho(\beta)} \leq 12 e^{\beta(V_{\text{max}}-V_{\text{min}})} e^{-\frac{2a^2}{\beta}}. \quad (2.22)$$

In the Appendix, a similar result is proved for the density matrix $\rho(\underline{r}, \underline{r}'; \beta)$.

3. Discussion

The key feature of Eq. (2.22) is the Gaussian decrease of $\Delta\rho(0, \beta)/\rho(0, \beta)$ with the size $2a$ of the cube which separates the interior and exterior regions. Thus we see that any change of the potential $V(\underline{r})$ outside of Ω_{int} , subject only to the lower bound limitation (2.7), will have a negligible effect on $\rho(0, \beta)$, provided that a is large enough. In particular, the potential outside of Ω_{int} may be replaced by zero or Ω_{int} may be enclosed by an infinite barrier, without having an appreciable effect on $\rho(0, \beta)$.

Since $\rho(\underline{r}, \beta)$ is the Laplace transform of the local density of states $n(\underline{r}, E)$ (See Eqs. (1.8) and (1.10)) the "locality" of $\rho(\underline{r}, \beta)$ implies in a general way the locality of $n(\underline{r}, E)$. Thus the theorem (2.22) which we have derived provides a general support for the physical fact that, in an extended system physical properties near a point \underline{r} depend significantly only on the potential near \underline{r} . It also lends support for various local methods of calculation, such as cluster methods⁽³⁻⁵⁾, moment methods⁽⁶⁻⁷⁾, etc., which have been extensively used in recent years to determine local properties of extended systems.

However, the Gaussian decay of $\Delta\rho(\underline{r}, \beta)$ as function of a , by no means implies that the inverse Laplace transform of $\rho(\underline{r}, \beta)$, the local density of

states $n(r, E)$, will also be local in the same very strong sense. $n(r, E)$ does in fact also obey a locality principle but in a much weaker sense, about which further details will be published in a contribution to the Festschrift for Felix Bloch (1979).

APPENDIX

In the following, we prove a localization theorem for the Laplace transform of the density matrix,

$$\rho(\underline{r}, \underline{r}'; \beta) \equiv \int n(\underline{r}, \underline{r}'; E) e^{-\beta E} dE, \quad (\text{A.1})$$

where $n(\underline{r}, \underline{r}'; E)$ is the density matrix of the system under consideration,

$$n(\underline{r}, \underline{r}'; E) = \sum_V \psi^{V*}(\underline{r}) \psi^V(\underline{r}') \delta(E - E^V). \quad (\text{A.2})$$

The Laplace transform $\rho(\underline{r}, \underline{r}'; \beta)$ can be written as a path integral similar to Eq. (2.1), where the relevant paths are those starting at \underline{r} and ending at \underline{r}' . Thus, for $\underline{r}, \underline{r}'$ inside Ω_{int} , $\rho(\underline{r}, \underline{r}'; \beta)$ obeys inequalities analogous to Eqs. (2.8) and (2.13). Consequently, in order to show that $\rho(\underline{r}, \underline{r}'; \beta)$ is localized, it suffices to show that $\rho_0(\underline{r}, \underline{r}'; \beta)$ has this property, where ρ_0 is the corresponding free-electron quantity.

The contribution of paths, entirely inside the cube Ω_{int} , to ρ_0 can be evaluated explicitly by using (A.1) and (A.2) and the appropriate wave functions. In this way we obtain

$$\rho_{0,\text{int}} = J(x, x') J(y, y') J(z, z'), \quad (\text{A.3})$$

where

$$J(x, x') = \sum_n \psi_n^*(x) \psi_n(x') e^{-\beta E_n}, \quad (\text{A.4})$$

and the set ψ_n is given by

$$\begin{aligned} \psi_{2n}(x) &= a^{-1/2} \cos(k_{2n} x) ; \quad k_{2n} a = \pi(n + 1/2) \\ \psi_{2n+1}(x) &= a^{-1/2} \sin(k_{2n+1} x) ; \quad k_{2n+1} a = \pi(n+1) \\ E_n &= \frac{1}{2} k_n^2 ; \quad n = 0, 1, 2, \dots \end{aligned}$$

Separating J into even and odd contributions, and using Poisson's summation formula, one can show that

$$J = \frac{e^{-\frac{(x-x')^2}{2\beta}}}{(2\pi\beta)^{1/2}} \left[1 + 2 \sum_{n=1}^{\infty} e^{-\frac{8a^2 n^2}{\beta}} \cosh\left[\frac{4an(x-x')}{\beta}\right] - 2 e^{-\frac{2xx'}{\beta}} \sum_{n=1}^{\infty} e^{-\frac{2a^2(2n-1)^2}{\beta}} \cosh\left[\frac{2a(2n-1)(x+x')}{\beta}\right] \right] \quad (\text{A.5})$$

For a large cube of edge $2a$, and for x and x' close enough to the origin (in such a way that $|x| + |x'| \ll a$), the series in (A.5) converge very rapidly and can be replaced by the corresponding first terms. In this way we obtain

$$J(x, x') = J_{\infty}(x, x') \left\{ 1 + 2e^{-\frac{8a^2}{\beta}} \cosh\left[\frac{4a(x-x')}{\beta}\right] - 2e^{-\frac{2(a^2+xx')}{\beta}} \cosh\left[\frac{2a(x+x')}{\beta}\right] \right\} \quad (\text{A.6})$$

where J_{∞} is the corresponding contribution from an infinite system, i.e.,

$$J_{\infty}(x, x') = (2\pi\beta)^{-1/2} e^{-\frac{(x-x')^2}{2\beta}} \quad (\text{A.7})$$

We apply now Eqs. (2.13), (A.3) and (A.8) to obtain the following localization theorem:

$$\frac{|\Delta\rho(\underline{r}, \underline{r}'; \beta)|}{\rho(\underline{r}; \underline{r}'; \beta)} \leq 2e^{\beta(V_{\max} - V_{\min})} \left[1 - g(x, x') g(y, y') g(z, z') \right], \quad (\text{A.8})$$

where $\Delta\rho$ is the change in ρ due to the change in the potential outside of Ω_{int} , and g is given by

$$g(x, x') = 1 - 2e^{-\frac{2a^2}{\beta}} \left\{ e^{-\frac{2xx'}{\beta}} \cosh\left[\frac{2a(x+x')}{\beta}\right] - e^{-\frac{6a^2}{\beta}} \cosh\left[\frac{4a(x-x')}{\beta}\right] \right\} \quad (\text{A.9})$$

We note that the relation (2.22) is a particular case of the general result (A.8) with $\underline{r} = \underline{r}' = 0$.

REFERENCES

1. Feynman R. P. and Hibbs A. R. (1965) Quantum Mechanics and Path Integrals (McGraw Hill, New York).
2. Bellman R. (1961) A Brief Introduction to Theta Functions (Holt, Rinehart and Winston, New York).
3. Kohn W. and Butler W. H. (1970) J. of Research of the National Bureau of Standards, Phys. and Chem. 74A, 443-447.
4. Keller J. (1972) J. de Physique, Supplement au N° 5-6 Tome 33, C3, 241-244.
5. Messmer R. P., Knudson S. K., Johnson K. H., Diamond J. B., and Yang C. Y. (1976) Phys. Rev. B13, 1396-1415.
6. Cyrot-Lackman F. (1968) J. Phys. Chem. Solids 29, 1235-1243.
7. Haydock R., Heine V., and Kelly M. J. (1975) J. Phys. C8, 2591-2605.